

# COMPSCI 514: ALGORITHMS FOR DATA SCIENCE

---

Cameron Musco

University of Massachusetts Amherst. Fall 2019.

Lecture 12

- Problem Set 2 is due this Friday 10/11. Will allow submissions until Sunday 10/13 at midnight with no penalty.
- No class next Tuesday (Monday class schedule). I will hold office hours from 10:30am-12:30 pm.

## **Midterm next Thursday 10/17 in class.**

- See review material posted with class schedule.
- More short-answer style than the problem sets.
- Review sheet will let you know what you need to memorize and what you don't.

### Last Class: Low-Rank Approximation and PCA


- How to compress a dataset that lies close to a  $k$ -dimensional subspace.
- View as projection, low-rank approximation of the data matrix  $\mathbf{X}$ .
- View as finding a small set of basis vectors for the rows or the columns of  $\mathbf{X}$ .

### This Class: Finish low-rank approximation and connection to eigendecomposition.

- Show how to find the best rank- $k$  subspace to approximation  $\mathbf{X}$  via eigendecomposition.
- Show how to calculate the error of the approximation.

## REVIEW OF LAST TIME

**Set Up:** Assume that data points  $\vec{x}_1, \dots, \vec{x}_n$  lie close to any  $k$ -dimensional subspace  $\mathcal{V}$  of  $\mathbb{R}^d$ . Let  $\mathbf{X} \in \mathbb{R}^{n \times d}$  be the data matrix.




Let  $\vec{v}_1, \dots, \vec{v}_k$  be an orthonormal basis for  $\mathcal{V}$  and  $\mathbf{V} \in \mathbb{R}^{d \times k}$  be the matrix with these vectors as its columns.

- $\mathbf{W}\mathbf{W}^T \in \mathbb{R}^{d \times d}$  is the **projection matrix** onto  $\mathcal{V}$ .
- $\mathbf{X} \approx \mathbf{X}(\mathbf{W}\mathbf{W}^T)$ . Gives the closest approximation to  $\mathbf{X}$  with rows in  $\mathcal{V}$ .

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : orthogonal basis for subspace  $\mathcal{V}$ .  $\mathbf{V} \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

## REVIEW OF LAST TIME


Low-Rank Approximation: Approximate  $X \approx XVV^T$ .



- $XVV^T$  is a **rank- $k$  matrix** – all its rows fall in  $\mathcal{V}$ .
- $X$ 's rows are approximately spanned by the columns of  $V$ .
- $X$ 's columns are approximately spanned by the columns of  $XV$ .

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $X \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : orthogonal basis for subspace  $\mathcal{V}$ .  $V \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

# DUAL VIEW OF LOW-RANK APPROXIMATION



Row (data point) compression

Column (feature) compression

$10000 * \text{bathrooms} + 10 * (\text{sq. ft.}) \approx \text{list price}$

	bedrooms	bathrooms	sq.ft.	floors	list price	sale price
home 1	2	2	1800	2	200,000	195,000
home 2	4	2.5	2700	1	300,000	310,000
.	.	.	.	.	.	.
.	.	.	.	.	.	.
.	.	.	.	.	.	.
home n	5	3.5	3600	3	450,000	450,000

## FINDING THE SUBSPACE

Given  $\vec{x}_1, \dots, \vec{x}_n$  that are close to a  $k$ -dimensional subspace  $\mathcal{V}$ ,

How do we find  $\mathcal{V}$  (and  $\mathbf{V}$ )?

$$\arg \min_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2 = \sum_{i,j} (\mathbf{x}_{i,j} - (\mathbf{X}\mathbf{V}\mathbf{V}^T)_{i,j})^2 = \sum_{i=1}^n \|\vec{x}_i - \mathbf{V}\mathbf{V}^T\vec{x}_i\|_2^2$$

By Pythagorean theorem, minimizing this error is the same as maximizing the norm of the projected dataset:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2 = \sum_{i=1}^n \|\mathbf{V}\mathbf{V}^T\vec{x}_i\|_2^2$$

Projection only reduces data point lengths and distances. Want to minimize this reduction. How does this compare to JL random projection?

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : orthogonal basis for subspace  $\mathcal{V}$ .  $\mathbf{V} \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

$\mathbf{V}$  minimizing  $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$  is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2 = \sum_{i=1}^n \|\mathbf{V}\mathbf{V}^T \vec{x}_i\|_2^2$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : orthogonal basis for subspace  $\mathcal{V}$ .  $\mathbf{V} \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .



$\mathbf{V}$  minimizing  $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$  is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2 = \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : orthogonal basis for subspace  $\mathcal{V}$ .  $\mathbf{V} \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

$\mathbf{V}$  minimizing  $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$  is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2 = \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_{j=1}^k \sum_{i=1}^n \langle \vec{v}_j, \vec{x}_i \rangle^2$$


$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : orthogonal basis for subspace  $\mathcal{V}$ .  $\mathbf{V} \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

## BEST FIT SUBSPACE

$\mathbf{V}$  minimizing  $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$  is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2 = \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_{j=1}^k \sum_{i=1}^n \langle \vec{v}_j, \vec{x}_i \rangle^2$$

Columns of  $\mathbf{V}$  are ‘directions of greatest variance’ in the data.




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : orthogonal basis for subspace  $\mathcal{V}$ .  $\mathbf{V} \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

$\mathbf{V}$  minimizing  $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$  is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2 = \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_{j=1}^k \sum_{i=1}^n \langle \vec{v}_j, \vec{x}_i \rangle^2$$

Columns of  $\mathbf{V}$  are ‘directions of greatest variance’ in the data.



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : orthogonal basis for subspace  $\mathcal{V}$ .  $\mathbf{V} \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

## SOLUTION VIA EIGENDECOMPOSITION

$\mathbf{V}$  minimizing  $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$  is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X}\mathbf{V}\|_F^2 = \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_{j=1}^k \sum_{i=1}^n \langle \vec{v}_j, \vec{x}_i \rangle^2 = \sum_{j=1}^k \|\mathbf{X}\vec{v}_j\|_2^2$$

Surprisingly, can find the columns of  $\mathbf{V}$ ,  $\vec{v}_1, \dots, \vec{v}_k$  **greedily!**

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

$$\vec{v}_2 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_1 \rangle=0} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

...

$$\vec{v}_k = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_j \rangle=0 \ \forall j < k} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

These are exactly the top  $k$  eigenvectors of  $\mathbf{X}^T \mathbf{X}$ .

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : orthogonal basis for subspace  $\mathcal{V}$ .  $\mathbf{V} \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .


**Eigenvector:**  $\vec{x} \in \mathbb{R}^d$  is an eigenvector of a matrix  $\mathbf{A} \in \mathbb{R}^{d \times d}$  if  $\mathbf{A}\vec{x} = \lambda\vec{x}$  for some scalar  $\lambda$  (the eigenvalue corresponding to  $\vec{x}$ ).

- That is,  $\mathbf{A}$  just ‘stretches’  $x$ .
- If  $\mathbf{A}$  is **symmetric**, can find  $d$  orthonormal eigenvectors  $\vec{v}_1, \dots, \vec{v}_d$ . Let  $\mathbf{V} \in \mathbb{R}^{d \times d}$  have these vectors as columns.

$$\mathbf{AV} = \begin{bmatrix} | & | & | & | \\ \mathbf{A}\vec{v}_1 & \mathbf{A}\vec{v}_2 & \cdots & \mathbf{A}\vec{v}_d \\ | & | & | & | \end{bmatrix} = \begin{bmatrix} | & | & | & | \\ \lambda_1\vec{v}_1 & \lambda_2\vec{v}_2 & \cdots & \lambda_d\vec{v}_d \\ | & | & | & | \end{bmatrix} = \mathbf{V}\mathbf{\Lambda}$$

Yields eigendecomposition:  $\mathbf{AVV}^T = \mathbf{A} = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^T$ .

# REVIEW OF EIGENVECTORS AND EIGENDECOMPOSITION



Typically order the eigenvectors in decreasing order:

$$\lambda_1 \geq \lambda_2 \geq \dots \lambda_d.$$

**Courant-Fischer Principal:** For symmetric  $\mathbf{A}$ , the eigenvectors are given via the greedy optimization:

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{A} \vec{v}.$$

$$\vec{v}_2 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_1 \rangle = 0} \vec{v}^T \mathbf{A} \vec{v}.$$


...

$$\vec{v}_d = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_j \rangle = 0 \ \forall j < d} \vec{v}^T \mathbf{A} \vec{v}.$$

- $\vec{v}_j^T \mathbf{A} \vec{v}_j = \lambda_j \cdot \vec{v}_j^T \vec{v}_j = \lambda_j$ , the  $j^{\text{th}}$  largest eigenvalue.
- The first  $k$  eigenvectors of  $\mathbf{X}^T \mathbf{X}$  (corresponding to the largest  $k$  eigenvalues) are exactly the directions of greatest variance in  $\mathbf{X}$  that we use for low-rank approximation.



# LOW-RANK APPROXIMATION VIA EIGENDECOMPOSITION



**Upshot:** Letting  $\mathbf{V}_k$  have columns  $\vec{v}_1, \dots, \vec{v}_k$  corresponding to the top  $k$  eigenvectors of the covariance matrix  $\mathbf{X}^T\mathbf{X}$ ,  $\mathbf{V}_k$  is the orthogonal basis minimizing

$$\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2,$$

This is principal component analysis (PCA).

How accurate is this low-rank approximation? Can understand using eigenvalues of  $\mathbf{X}^T\mathbf{X}$ .

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T\mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

Let  $\vec{v}_1, \dots, \vec{v}_k$  be the top  $k$  eigenvectors of  $\mathbf{X}^T \mathbf{X}$  (the top  $k$  principal components). Approximation error is:

$$\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 = \|\mathbf{X}\|_F^2 - \|\mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T \mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

Let  $\vec{v}_1, \dots, \vec{v}_k$  be the top  $k$  eigenvectors of  $\mathbf{X}^T \mathbf{X}$  (the top  $k$  principal components). Approximation error is:

$$\|\mathbf{X} - \mathbf{XV}_k \mathbf{V}_k^T\|_F^2 = \|\mathbf{X}\|_F^2 - \|\mathbf{XV}_k\|_F^2$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T \mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

Let  $\vec{v}_1, \dots, \vec{v}_k$  be the top  $k$  eigenvectors of  $\mathbf{X}^T \mathbf{X}$  (the top  $k$  principal components). Approximation error is:

$$\|\mathbf{X} - \mathbf{XV}_k \mathbf{V}_k^T\|_F^2 = \|\mathbf{X}\|_F^2 - \|\mathbf{XV}_k\|_F^2$$

- For any matrix  $\mathbf{A}$ ,  $\|\mathbf{A}\|_F^2 = \sum_{i=1}^d \|\vec{a}_i\|_2^2 = \text{tr}(\mathbf{A}^T \mathbf{A})$  (sum of diagonal entries = sum eigenvalues).

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T \mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

Let  $\vec{v}_1, \dots, \vec{v}_k$  be the top  $k$  eigenvectors of  $\mathbf{X}^T \mathbf{X}$  (the top  $k$  principal components). Approximation error is:


$$\begin{aligned} \|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 &= \text{tr}(\mathbf{X}^T \mathbf{X}) - \text{tr}(\mathbf{V}_k^T \mathbf{X}^T \mathbf{X} \mathbf{V}_k) \\ &= \sum_{i=1}^d \lambda_i(\mathbf{X}^T \mathbf{X}) - \sum_{i=1}^k \vec{v}_i^T \mathbf{X}^T \mathbf{X} \vec{v}_i \\ &= \sum_{i=1}^d \lambda_i(\mathbf{X}^T \mathbf{X}) - \sum_{i=1}^k \lambda_i(\mathbf{X}^T \mathbf{X}) = \sum_{i=k+1}^d \lambda_i(\mathbf{X}^T \mathbf{X}) \end{aligned}$$

- For any matrix  $\mathbf{A}$ ,  $\|\mathbf{A}\|_F^2 = \sum_{i=1}^d \|\vec{a}_i\|_2^2 = \text{tr}(\mathbf{A}^T \mathbf{A})$  (sum of diagonal entries = sum eigenvalues).

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T \mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

**Claim:** The error in approximating  $\mathbf{X}$  with the best rank  $k$  approximation (projecting onto the top  $k$  eigenvectors of  $\mathbf{X}^T\mathbf{X}$  is:

$$\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 = \sum_{i=k+1}^d \lambda_i(\mathbf{X}^T\mathbf{X})$$



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T\mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .


**Claim:** The error in approximating  $\mathbf{X}$  with the best rank  $k$  approximation (projecting onto the top  $k$  eigenvectors of  $\mathbf{X}^T\mathbf{X}$  is:

$$\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 = \sum_{i=k+1}^d \lambda_i(\mathbf{X}^T\mathbf{X})$$

784 dimensional vectors



eigendecomposition




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T\mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .




**Claim:** The error in approximating  $\mathbf{X}$  with the best rank  $k$  approximation (projecting onto the top  $k$  eigenvectors of  $\mathbf{X}^T\mathbf{X}$  is:

$$\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 = \sum_{i=k+1}^d \lambda_i(\mathbf{X}^T\mathbf{X})$$

784 dimensional vectors



eigendecomposition



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T\mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .


# SPECTRUM ANALYSIS

Plotting the **spectrum** of the covariance matrix  $\mathbf{X}^T\mathbf{X}$  (its eigenvalues) shows how compressible  $\mathbf{X}$  is using low-rank approximation (i.e., how close  $\vec{x}_1, \dots, \vec{x}_n$  are to a low-dimensional subspace).

784 dimensional vectors



eigendecomposition



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T\mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .


# SPECTRUM ANALYSIS

Plotting the **spectrum** of the covariance matrix  $\mathbf{X}^T\mathbf{X}$  (its eigenvalues) shows how compressible  $\mathbf{X}$  is using low-rank approximation (i.e., how close  $\vec{x}_1, \dots, \vec{x}_n$  are to a low-dimensional subspace).

784 dimensional vectors





eigendecomposition



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T\mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

# SPECTRUM ANALYSIS

Plotting the **spectrum** of the covariance matrix  $\mathbf{X}^T\mathbf{X}$  (its eigenvalues) shows how compressible  $\mathbf{X}$  is using low-rank approximation (i.e., how close  $\vec{x}_1, \dots, \vec{x}_n$  are to a low-dimensional subspace).



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T\mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

# INTERPRETATION IN TERMS OF CORRELATION

**Recall:** Low-rank approximation is possible when our data features are correlated.

10000\* bathrooms+ 10\* (sq. ft.)  $\approx$  list price

	bedrooms	bathrooms	sq.ft.	floors	list price	sale price
home 1	2	2	1800	2	200,000	195,000
home 2	4	2.5	2700	1	300,000	310,000
.	.	.	.	.	.	.
.	.	.	.	.	.	.
.	.	.	.	.	.	.
home n	5	3.5	3600	3	450,000	450,000

Our compressed dataset is  $\mathbf{C} = \mathbf{X}\mathbf{V}_k$  where the columns of  $\mathbf{V}_k$  are the top  $k$  eigenvectors of  $\mathbf{X}^T\mathbf{X}$ .

What is the covariance of  $\mathbf{C}$ ?  $\mathbf{C}^T\mathbf{C} = \mathbf{V}_k^T\mathbf{X}^T\mathbf{X}\mathbf{V}_k = \mathbf{V}_k^T\mathbf{V}\mathbf{V}^T\mathbf{V}_k = \mathbf{\Lambda}_k$

Covariance becomes diagonal. I.e., all correlations have been removed. Maximal compression.

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T\mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

What is the runtime to compute an optimal low-rank approximation?

- Computing the covariance matrix  $\mathbf{X}^T\mathbf{X}$  requires  $O(nd^2)$  time.
- Computing its full eigendecomposition to obtain  $\vec{v}_1, \dots, \vec{v}_k$  requires  $O(d^3)$  time (similar to the inverse  $(\mathbf{X}^T\mathbf{X})^{-1}$ ).

Many faster iterative and randomized methods. Runtime is roughly  $\tilde{O}(ndk)$  to output just the top  $k$  eigenvectors  $\vec{v}_1, \dots, \vec{v}_k$ .

- Will see in a few classes

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ : data points,  $\mathbf{X} \in \mathbb{R}^{n \times d}$ : data matrix,  $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$ : top eigenvectors of  $\mathbf{X}^T\mathbf{X}$ ,  $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ : matrix with columns  $\vec{v}_1, \dots, \vec{v}_k$ .

Questions?